



General Requisition

Date Specimen Collected: Time Specimen Collected: STAT

Laboratory Use Only

Accession Number Date Received Time Received

Practice Name Practice ID Practice Contact Information
Address
City, State, Zip
Phone
Fax

Patient and Insurance Information

(Currently Not Accepting Medicaid or Managed Medicaid Plans - Contract Pending)

First Name Last Name Middle Initial Gender*
Address Line 1 Address Line 2 City State Zip
DOB Cell/Home Phone* Email
Gender Identity* Race* Ethnicity* Sexual Orientation*
Insured's Name Relationship to Patient Social Security #
Home Phone Cell Phone DOB Gender
Primary Insurance Secondary Insurance
Group # ID # Group # ID#
Address Address
City State Zip City State Zip

Stool Test

Genesis Diarrhea Pathogen Panel (11 Pathogens with Genotypic Antibiotic Resistance)

Duration of Diarrhea:

> 7 days diarrhea
Other diarrhea duration

Urinary Tests

Urinary Tract Infection/Urinary Tract Microbiota (UTI/UTM) & Antibiotic Resistance Panel
Sexually Transmitted Infection (STI) Panel
Urinalysis (if abnormal, reflex UTI/UTM Panel)
Microalbumin
(beta)beta-hCG (pregnancy test)

Respiratory Tests

Nasal Swab OR Nasopharyngeal Swab:

Genesis Respiratory Pathogen Panel (Including COVID-19) (UTM Swab)
COVID-19 + Influenza A & B, and RSV Panel (UTM Swab)
COVID-19 ONLY (SARS-CoV-2 PCR Assay) (UTM Swab)
Genesis Expanded Respiratory Pathogen Panel** (UTM Swab)

Please note that an immunodeficiency diagnosis code must be selected in order for this test to be performed. If an appropriate immunodeficiency code is not selected, testing will automatically be reflexed to the "Genesis Respiratory Pathogen Panel".

Throat Swab:

Group A Streptococcus (Liquid Amies/eSwab Only)

Wound Panel

Wound Microbiota and Bacterial Genotypic Antibiotic Resistance by PCR with Gram Stain [Liquid Amies eSwab/Puritan Opti-Swab]

ICD-10 Codes

- R19.7 Diarrhea unspecified
R19.7/R50.9 Diarrhea, unspecified with fever
R19.7/K92.1 Diarrhea, unspecified with hematochezia
R19.7/R10.9 Diarrhea, unspecified with abdominal pain unspecified
R19.7/E87.8 Diarrhea, unspecified with other disorders of electrolyte and fluid imbalance
B95.61 Methicillin susceptible staphylococcus aureus infection
B95.62 Methicillin resistant staphylococcus aureus infection
N30.00 Acute cystitis
N34.1 Nonspecific urethritis
N34.2 Other urethritis
N39.0 Urinary Tract Infection
N91.2 Amenorrhea, unspecified
R30.0 Dysuria
Z11.3 Screen STD
R05 Cough
R50.9 Fever, unspecified
R68.83 Chills (without fever)
R59.9 Enlarged lymph nodes, unspecified
J06.9 Acute respiratory infection, unspecified
Z20.828 Contact with and (suspected) exposure to other viral communicable disease
L08.89 Other specified local infections of the skin and subcutaneous tissue
S91.301A Unspecified open wound, right foot, initial encounter
S91.302A Unspecified open wound, left foot, initial encounter
S91.102A Unspecified open wound of left great toe without damage to nail, initial encounter
S91.101A Unspecified open wound of right great toe without damage to nail, initial encounter
Immunodeficiency ICD-10 Codes
D84.821 Immunodeficiency due to drugs
D81.89 Other combined immunodeficiencies
D81.9 Combined immunodeficiency, unspecified
D83.8 Other common variable immunodeficiencies
D83.9 Common variable immunodeficiency, unspecified
D84.89 Other immunodeficiencies
D84.9 Immunodeficiency, unspecified

This test is medically necessary for the diagnosis or detection of a disease, illness, impairment, symptom, syndrome or disorder. The results will determine my patient's medical management and treatment decisions. The person listed as the ordering provider is authorized by law to order the test(s) requested herein.

Signature of Physician or Other Authorized NPI Provider (REQUIRED) Date Accessioner Initials 1 2

*Gender, Cell/Home Phone, Gender Identity, Race, Ethnicity, and Sexual Orientation are required by certain states and/or the CDC. See reverse side for panel details. ICD-10 Codes are listed for information purposes only. It is the provider's responsibility to order tests that are medically necessary and in the best interest of the patient.

Genesis Diarrhea Pathogen Panel and Genotypic Antibiotic Resistance Panel

<p>Bacteria <i>Campylobacter</i> (<i>C. jejuni</i> /<i>C. coli</i> /<i>C. upsaliensis</i>) <i>Clostridium difficile</i> (Toxin A/B)*** <i>Salmonella</i> spp. <i>Vibrio</i> spp. (<i>V. parahaemolyticus</i> /<i>V. vulnificus</i> / <i>V. cholerae</i>) <i>Yersinia enterocolitica</i></p>	<p>Viruses Norovirus GI/GII Rotavirus A, B, C</p>	<p>Bacterial Genotypic Antibiotic Resistance <i>Vancomycin Resistance</i>: <i>vanA</i>, <i>vanB</i>, <i>vanC</i> <i>Nitroimidazole Resistance</i>: <i>nimA</i>, <i>nimD</i> • <i>Examples</i>: <i>metronidazole</i>, <i>tinidazole</i> <i>Erythromycin Resistance</i>: <i>ermA</i>, <i>ermB</i>, <i>ermC</i>, <i>mefA</i> <i>Quinolone and Fluoroquinolone Resistance</i>: <i>qnrA</i>, <i>qnrB</i>, <i>qnrS</i> • <i>Examples</i>: <i>ciprofloxacin</i>, <i>levofloxacin</i>, <i>moxifloxacin</i> <i>Trimethoprim-Sulfamethoxazole resistance</i>: <i>dfrA</i>, <i>dfrA1</i>, <i>dfrA5</i></p>
<p>Diarrheagenic <i>E. coli</i> / <i>Shigella</i> Shiga-like toxin-producing <i>E. coli</i> (STEC) <i>Shigella</i> / Enteroinvasive <i>E. coli</i> (EIEC)</p>	<p>Parasites <i>Entamoeba histolytica</i> <i>Giardia lamblia</i></p>	

Respiratory Tests

<p>Genesis Respiratory Pathogen Panel</p> <p>Viruses Adenovirus COVID-19 (SARS-CoV-2) Human Metapneumovirus Influenza A virus / Influenza B virus Pan Respiratory syncytial virus</p> <p>Bacteria <i>Bordetella pan</i> (<i>bronchiseptica</i>, <i>parapertussis</i>, <i>pertussis</i>) <i>Bordetella pertussis</i> <i>Chlamydia pneumoniae</i> <i>Mycoplasma pneumoniae</i> <i>Legionella pneumophila</i></p>	<p>COVID-19 + Influenza A & B, and RSV Panel COVID-19 (SARS-CoV-2) Influenza A Influenza B Respiratory Syncytial Virus (RSV)</p> <p>COVID-19 ONLY COVID-19 (SARS-CoV-2)</p> <p>Group A Streptococcus <i>Streptococcus pyogenes</i></p>	<p>Genesis Expanded Respiratory Pathogen Panel</p> <p>Viruses Adenovirus Coronavirus 229E Coronavirus HKU1 Coronavirus NL63 Coronavirus OC43 COVID-19 (SARS-CoV-2) Human Metapneumovirus Human Rhinovirus / Enterovirus Influenza A virus Influenza A virus A/H1 Influenza A virus A/H3 Influenza A virus A/H1-2009 Influenza B virus Parainfluenza virus 1</p> <p>Parainfluenza virus 2 Parainfluenza virus 3 Parainfluenza virus 4 Respiratory syncytial virus</p> <p>Bacteria <i>Bordetella pan</i> (<i>bronchiseptica</i>, <i>parapertussis</i>, <i>pertussis</i>) <i>Bordetella pertussis</i> <i>Chlamydia pneumoniae</i> <i>Mycoplasma pneumoniae</i></p>
--	---	---

Urinary Tests

<p>Urinary Tract Infection / Urinary Tract Microbiota Panel</p> <p>Gram-negative microbiota <i>Escherichia coli</i> <i>Klebsiella pneumoniae</i> <i>Proteus mirabilis</i> <i>Pseudomonas aeruginosa</i> <i>Providencia stuartii</i> <i>Morganella morganii</i></p>	<p>Gram-positive microbiota <i>Enterococcus faecalis</i> <i>Enterococcus faecium</i> <i>Streptococcus agalactiae</i> <i>Staphylococcus saprophyticus</i></p> <p>Fungal microbiota <i>Candida albicans</i></p>	<p>Urinary STI Tests <i>Chlamydia trachomatis</i> (CT) <i>Neisseria gonorrhoeae</i> (NG) <i>Trichomonas vaginalis</i> (TV) <i>Mycoplasma genitalium</i> (MG)</p>
--	--	---

Wound Microbiota and Bacterial Genotypic Antibiotic Resistance by PCR with Gram Stain

<p>Gram-Negative Bacteria <i>Acinetobacter baumannii</i> <i>Bacteroides fragilis</i> <i>Bacteroides vulgatus</i> <i>Citrobacter freundii</i> <i>Escherichia coli</i> <i>Enterobacter cloacae</i> <i>Fusobacterium necrophorum</i> <i>Klebsiella pneumoniae</i> <i>Proteus mirabilis</i> <i>Pseudomonas aeruginosa</i> <i>Morganella morganii</i></p>	<p>Gram-Positive Bacteria <i>Clostridium perfringens</i> <i>Clostridium septicum</i> <i>Corynebacterium striatum</i> <i>Enterococcus faecalis</i> <i>Finegoldia magna</i> <i>Peptoniphilus harei</i> <i>Peptostreptococcus anaerobius</i> <i>Peptostreptococcus asaccharolyticus</i> <i>Peptostreptococcus prevotii</i> <i>Staphylococcus aureus</i> <i>Staphylococcus haemolyticus</i> <i>Staphylococcus lugdunensis</i> <i>Staphylococcus saprophyticus</i></p>	<p>Streptococcus agalactiae <i>Streptococcus pyogenes</i></p>	<p>Genotypic Resistance</p> <p><i>ampC</i> <i>KPC</i> <i>OXA-1</i> <i>NDM</i> <i>CTX-M Group2</i> <i>ErmC</i> <i>mecC</i> <i>vanA</i></p> <p><i>VIM</i> <i>IMP-7</i> <i>OXA-48</i> <i>CTX-M Group1</i> <i>ErmB</i> <i>mecA</i> <i>QnrA</i> <i>vanB</i></p>
			<p>Fungal <i>Candida albicans</i> <i>Candida auris</i> <i>Candida parapsilosis</i></p>

***Reflex to active Toxin detection by EIA for positive samples